



Join us for a free NF1 PN educational virtual event.



Managing NF1 PN can be complex—that's why **KOSELUGO® (selumetinib) comes with a team.**

At Alexion, we know that support matters. That's why Koselugo comes with a team. Hear from a Patient Education Manager and a healthcare provider to learn more about neurofibromatosis type 1 with plexiform neurofibromas (NF1 PN).

At this event you will learn more about:

- NF1 PN and a treatment option
- How Koselugo can help
- Support for families taking Koselugo
- NF1 and NF1 PN
- Next steps for parents and caregivers
- Alexion's support available for NF1 families

For more information about Programs, reach out to a Patient Education Manager (PEM)

To find the PEM closest to you, use the PEM finder at bit.ly/NF1-PEM-Finder.

INDICATION

What is Koselugo?

Koselugo is a prescription medicine that is used to treat children 2 years of age and older with neurofibromatosis type 1 (NF1) who have plexiform neurofibromas that cannot be completely removed by surgery. It is not known if Koselugo is safe and effective in children under 2 years of age.

Please see full Important Safety Information on the next page and [click here](#) to see accompanying full Prescribing Information for Koselugo.

TOPIC:

All about NF1, PN, and a treatment option

WHO:

Breanne Roche
CNP, DNP

WHEN:

Thursday, April 25, 2024
7:00 PM EST

WHERE:

Virtual

To register and/or learn more about events, please scan the QR code, call 888-210-0174, or visit koselugo.com/events.



IMPORTANT SAFETY INFORMATION

What are the possible side effects of Koselugo?

Koselugo may cause serious side effects, including:

- **Heart problems.** Koselugo can lower the amount of blood pumped by your heart, which is common and can also be severe. Your healthcare provider will do tests before and during treatment with Koselugo to check how well your heart is working. Tell your healthcare provider right away if you get any of the following signs or symptoms: persistent coughing or wheezing, shortness of breath, swelling of your ankles and feet, tiredness, increased heart rate.
- **Eye problems.** Koselugo can cause eye problems that can lead to blindness. Your healthcare provider will check your vision before and during treatment with Koselugo. Tell your healthcare provider right away if you get any of the following signs or symptoms: blurred vision, loss of vision, dark spots in your vision (floaters), other changes to your vision.
- **Severe diarrhea.** Diarrhea is common with Koselugo and can also be severe. Tell your healthcare provider right away the first time that you get diarrhea during treatment with Koselugo. Your healthcare provider may give you medicine to help control your diarrhea and may tell you to drink more fluids.
- **Skin rash.** Skin rashes are common with Koselugo and can also be severe. Tell your healthcare provider if you get any of the following signs or symptoms: rash that covers a large area of your body, peeling skin, blisters.
- **Muscle problems (rhabdomyolysis).** Muscle problems are common with Koselugo and can also be severe. Treatment with Koselugo may increase the level of a muscle enzyme in your blood called creatine phosphokinase (CPK) and may be a sign of muscle damage. Your healthcare provider should do a blood test to check your muscle enzyme levels of CPK before you start taking Koselugo and during treatment. Tell your healthcare provider right away if you get any of the following signs or symptoms: muscle aches or pain, muscle spasms and weakness, dark, reddish urine.

Before taking Koselugo, tell your healthcare provider about all your medical conditions, including if you:

- have heart problems.
- have eye problems.
- are pregnant or plan to become pregnant. Koselugo can harm your unborn baby.
 - Your healthcare provider should verify if you/your partner are pregnant before beginning Koselugo.
 - Ensure you/your partner use effective birth control (contraception) during treatment with Koselugo and for 1 week after your last dose if there is possibility pregnancy could occur.
 - Tell your healthcare provider right away if you/your partner become pregnant or think you/your partner may be pregnant during treatment with Koselugo.
- are breastfeeding or plan to breastfeed. It is not known if Koselugo passes into your breast milk.
 - Do not breastfeed during treatment with Koselugo and for 1 week after your last dose.
 - Talk to your healthcare provider about the best way to feed your baby during this time.

The most common side effects of Koselugo are: vomiting, stomach pain, nausea, dry skin, feeling of tiredness, weakness or lacking energy, muscle and bone pain, fever, inflammation of the mouth, headache, redness around the fingernails, itching.

Tell your healthcare provider about all the medicines you take, including prescription, over-the-counter medicines, vitamins, or herbal supplements. Especially tell your healthcare provider if you are taking aspirin, blood thinners, or other medicines to treat blood clots. Koselugo contains vitamin E, which may increase risk of bleeding.

What should I avoid while taking Koselugo?

Do not drink grapefruit juice, eat grapefruit, or take supplements with grapefruit or St. John's Wort during treatment.

These are not all the possible side effects of Koselugo. Call your doctor for medical advice about side effects. Your healthcare provider may change your dose, temporarily stop, or permanently ask you to stop taking Koselugo if you have any of these side effects. **You may report side effects to AstraZeneca at 1-800-236-9933 or at <https://us-aereporting.astrazeneca.com> or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.**

Please see full Important Safety Information above or [click here](#) to see accompanying full Prescribing Information for Koselugo.