



KOSELUGO® (selumetinib) is a prescription medicine that is used to treat children 2 years of age and older with neurofibromatosis type 1 (NF1) who have plexiform neurofibromas that cannot be completely removed by surgery. It is not known if Koselugo is safe and effective in children under 2 years of age.

Taking Koselugo just got more convenient



Caleb, age 7, living with NF1 PN. Caleb is not a Koselugo patient.

Koselugo can be taken on an empty stomach or with food.

Enjoy more flexibility in your daily life without having to follow a fasting schedule.

 Consult with your loved one's doctor about taking Koselugo with food.

 Take Koselugo at a time that works for you and your loved one, twice daily, about 12 hours apart.

 Koselugo capsules must only be swallowed whole with water. They must not be chewed, dissolved, or opened.



Actor portrayal

Have questions?

Your dedicated local Patient Education Manager (PEM) is available to meet in person or virtually to help.

[Contact PEM](#)

IMPORTANT SAFETY INFORMATION for KOSELUGO® (selumetinib)

What are the possible side effects of Koselugo?

Koselugo may cause serious side effects, including:

- **Heart problems.** Koselugo can lower the amount of blood pumped by your heart, which is common and can also be severe. Your healthcare provider will do tests before and during treatment to check how well your heart is working. Tell your healthcare provider right away if you get any of the following signs or symptoms: persistent coughing or wheezing, shortness of breath, swelling of your ankles and feet, tiredness, increased heart rate.
- **Eye problems.** Koselugo can cause eye problems that can lead to blindness. Your healthcare provider will check your vision before and during treatment. Tell your healthcare provider right away if you get any of the following signs or symptoms: blurred vision, loss of vision, dark spots in your vision (floaters), other changes to your vision.
- **Severe diarrhea.** Diarrhea is common with Koselugo and can also be severe. Tell your healthcare provider right away the first time that you get diarrhea during treatment. Your healthcare provider may give you medicine to help control your diarrhea and may tell you to drink more fluids.
- **Skin rash.** Skin rashes are common with Koselugo and can also be severe. Tell your healthcare provider if you get any of the following signs or symptoms: rash that covers a large area of your body, peeling skin, blisters.
- **Muscle problems (rhabdomyolysis).** Muscle problems are common with Koselugo and can also be severe. Treatment with Koselugo may increase the level of a muscle enzyme in your blood called creatine phosphokinase (CPK) and may be a sign of muscle damage. Your healthcare provider should do a blood test to check your muscle enzyme levels of CPK before you start taking Koselugo and during treatment. Tell your healthcare provider right away if you get any of the following signs or symptoms: muscle aches or pain; muscle spasms and weakness; dark, reddish urine.

Before taking Koselugo, tell your healthcare provider about all your medical conditions, including if you:

- have heart problems.
- have eye problems.
- have liver problems.
- are pregnant or plan to become pregnant. Koselugo can harm your unborn baby. Your healthcare provider should verify if you/your partner are pregnant before beginning treatment. Ensure you/your partner use effective birth control (contraception) during treatment and for 1 week after your last dose if there is possibility pregnancy could occur. Tell your healthcare provider right away if you/your partner think you may be pregnant.
- are breastfeeding or plan to breastfeed. It is not known if Koselugo passes into your breast milk. Do not breastfeed during treatment and for 1 week after your last dose. Talk to your healthcare provider about the best way to feed your baby during this time.

Tell your healthcare provider about all the medicines you take, including prescription, over-the-counter medicines, vitamins, or herbal supplements. Especially tell your healthcare provider if you are taking aspirin, blood thinners, or other medicines to treat blood clots. Koselugo contains vitamin E, which may increase risk of bleeding.

What should I avoid while taking Koselugo?

Do not drink grapefruit juice, eat grapefruit, or take supplements with grapefruit or St. John's Wort during treatment.

Most common side effects include: vomiting, stomach-area pain, nausea, dry skin, muscle and bone pain, feeling of tiredness or lacking energy, fever, sores in your mouth, headache, redness around the fingernails, itching.

These are not all the possible side effects of Koselugo. Call your healthcare provider for medical advice about side effects. Your healthcare provider may change your dose, temporarily stop, or permanently ask you to stop taking Koselugo if you have any of these side effects. **You may report side effects to AstraZeneca at 1-800-236-9933 or at <https://us-aereporting.astrazeneca.com> or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.**

Please see [Patient Information](#) in the full [Prescribing Information](#) for Koselugo (selumetinib).

Caleb and his family have been compensated by Alexion.

[Contact Us](#) [Unsubscribe](#) [Privacy Policy](#)

This email was sent by **Alexion Pharmaceuticals, Inc.**
121 Seaport Blvd - Boston, MA 02210 - United States



KOSELUGO is a registered trademark of the AstraZeneca group of companies. ALEXION and the Alexion logo are registered trademarks of Alexion Pharmaceuticals, Inc.

© 2024, Alexion Pharmaceuticals, Inc. All rights reserved.

US/KOS-NF1/0631 V1 02/2024